

Henry Blackburn,

3 September 1988

You may be interested in what is below. Should a table like this be added to the ms "Fatness and Longevity?"

The Framingham Study, men 55-64, reported by D. Shurtleff in "Some characteristics related to the incidence of cardiovascular disease and death, 18-year follow-up. Epidemiological investigation of cardiovascular disease, 1974, Section 30 DHEW no. (NIH) 74-599.

Below are the multiple logistic regression coefficients (COEFF), their standard errors, and t-values for 1) the regression of death rate on the characteristic (Univariate), 2) the regression of death rate on the characteristic taking into account age, (Bivariate), 3) the regression of death rate taking into account age, systolic blood pressure, serum cholesterol, cigarette use, glucose intolerance, left ventricular hypertrophy (ECG).

CHARACTERISTIC	REGRESSION	COEFF	S.E.	T-VALUE
Systolic B P	Univariate	0.0160	0.0018	8.87
	Bivariate	0.0117	0.0019	6.29
	Multivar.	0.0054	0.0021	2.64
Cigarettes	Univariate	0.0972	0.0404	2.41
	Bivariate	0.2224	0.0421	5.29
	Multivar.	0.2638	0.0433	6.09
Relative wt. (BMI)	Univariate	-0.0134	0.0031	-4.28
	Bivariate	-0.0117	0.0032	-3.72
	Multivar.	-0.0149	0.0033	-4.48

Ancel Keys in residence:

15 September - 15 December, "Minnelea" 84060 Pioppi (SA) Italy

16 December - 15 January 1989, 410 Groveland Av. (No. 802)

Minneapolis Minn 55403, USA.